

TWISTED CONJUGACY AND SEPARABILITY

SAM TERTOORY 

ABSTRACT. A group G is twisted conjugacy separable if for every automorphism φ , distinct φ -twisted conjugacy classes can be separated in a finite quotient. Likewise, G is completely twisted conjugacy separable if for any group H and any two homomorphisms φ, ψ from H to G , distinct (φ, ψ) -twisted conjugacy classes can be separated in a finite quotient. We study how these properties behave with respect to taking subgroups, quotients and finite extensions, and compare them to other notions of separability in groups. Finally, we show that for polycyclic-by-nilpotent-by-finite groups, being completely twisted conjugacy separable is equivalent to all quotients being residually finite.

1. INTRODUCTION

Let G and H be groups and let $\varphi, \psi: H \rightarrow G$ be two homomorphisms. Two elements g_1, g_2 of G are said to be (φ, ψ) -twisted conjugate if there exists some $h \in H$ such that $g_1 = \psi(h)g_2\varphi(h)^{-1}$. This creates an equivalence relation, the classes of which are called (φ, ψ) -twisted conjugacy classes; the class of $g \in G$ is denoted by $[g]_{\varphi, \psi}$. If $G = H$ and $\psi = \text{id}_G$, we omit ψ and just write ‘ φ -twisted conjugacy’ and $[g]_{\varphi}$. If moreover $\varphi = \text{id}_G$ as well, we have the usual conjugacy relation and we write $[g]$ for the conjugacy class of $g \in G$.

In [24], Stebe introduced the notion of nests in a group G . A subset of $G \times G$ is called a *pre-nest* if it contains $(ac^{-1}, d^{-1}b)$ whenever it contains both (a, b) and (c, d) . The set of all elements $ab \in G$ corresponding to the elements (a, b) of a pre-nest is called a *nest* in G . Examples of nests include subgroups, products of two subgroups and twisted conjugacy classes of the identity.

A subset S of a group G is called *separable* if for every $g \notin S$, there is a finite index normal subgroup N of G such that $g \notin SN$. This gives rise to many ‘separability properties’, e.g. a group G is called *residually finite* if all singletons $\{g\}$ are separable, *strongly residually finite* if all normal subgroups are separable, *extended residually finite* if all subgroups are separable, *conjugacy separable* if all conjugacy classes are separable and *residually finite with respect to nests* if all nests are separable.

In [2], Deré and Pengitore defined a group G to be *twisted conjugacy separable* if for all $\varphi \in \text{Aut}(G)$ and all $g \in G$, the twisted conjugacy class $[g]_{\varphi}$ is separable. It should be noted that the term ‘twisted conjugacy separable’ first occurred in the works of Fel’shtyn and Troitsky [4, 5]. However, their definition differs from the one given by Deré and Pengitore, which is the one we shall use.

In this paper, we introduce and study *complete twisted conjugacy separability*. A group G is said to have this property if for all groups H , all $\varphi, \psi \in \text{Hom}(H, G)$ and all $g \in G$, the twisted conjugacy class $[g]_{\varphi, \psi}$ is separable. Unlike (twisted) conjugacy separability, this property is retained when taking subgroups, quotients and finite extensions. Our main results are the following.

Date: 29th February 2024.

2020 Mathematics Subject Classification. Primary: 20E26; Secondary: 20E18, 20E45, 20F19.

Key words and phrases. Twisted conjugacy, separability, residual finiteness, nests, nilpotent groups, polycyclic groups, profinite groups.

Theorem A. *A group is residually finite with respect to nests if and only if it is completely twisted conjugacy separable.*

Theorem B. *For a polycyclic-by-nilpotent-by-finite group G , the following are equivalent:*

- (a) G is strongly residually finite;
- (b) G is extended residually finite;
- (c) G is residually finite with respect to nests;
- (d) G is completely twisted conjugacy separable.

Theorem A shows that residual finiteness with respect to nests, or even nests in general, can be studied using twisted conjugacy as a framework. Theorem B extends a result by Menth [15], who proved that strong and extended residual finiteness are equivalent for nilpotent-by-finite groups.

We finish this section by agreeing on some notation. For a group G , we will always use \bar{G} and \tilde{G} to denote quotients of G . For an element g , we use \bar{g} and \tilde{g} to denote the projections of g to \bar{G} and \tilde{G} respectively. For a homomorphism $\varphi \in \text{Hom}(H, G)$, whenever we write $\bar{\varphi} \in \text{Hom}(H, \bar{G})$, we mean the homomorphism induced by φ , i.e. $\bar{\varphi} := p \circ \varphi$ with $p: G \rightarrow \bar{G}$ the quotient map. We will also encounter \hat{G} , however, the ‘hat’ will always indicate the profinite completion of G and never a quotient of G . We will use $H \leq_f G$ and $N \trianglelefteq_f G$ to indicate a finite index subgroup H and a finite index normal subgroup N respectively. The inner automorphism of G that conjugates every element by $g \in G$, will be denoted by ι_g .

2. SEPARABILITY PROPERTIES

A subset S of a group G is called *separable* if any of the following equivalent conditions hold:

- For each $g \in G \setminus S$, there exists $N \trianglelefteq_f G$ such that $g \notin SN$;
- For each $g \in G \setminus S$, there exists a homomorphism p from G onto a finite group such that $p(g) \notin p(S)$;
- S is closed in the finite-index topology of G .

The finite-index topology mentioned here is the topology on a group obtained by taking as a basis the set of all cosets (both left and right) of all finite index subgroups. Any group becomes a topological group when equipped with this topology. This topology is often called the ‘‘profinite topology’’, however, as we will be dealing with profinite groups and their topology later in this paper, we avoid this name to prevent any potential confusion between the two.

We now give an overview of various separability properties. We give their definition, mention whether they are retained or not under taking subgroups, quotients and finite extensions, and discuss how they are related to one another. Only results available in the literature are considered in this section.

Definition 2.1. A group G is called *residually finite (RF)* if every singleton $\{g\} \subseteq G$ is separable.

There are many equivalent ways to define residual finiteness. It suffices to demand that $\{1_G\}$ is separable, or, one can demand that every finite subset of G is separable. From a topological point of view, G is residually finite if and only if its finite-index topology is Hausdorff.

The family of RF-groups is subgroup-closed and closed under taking finite extensions, but not quotient-closed. For example, the group of dyadic rationals $\mathbb{Z}[\frac{1}{2}]$ is residually finite, yet its quotient $\mathbb{Z}[\frac{1}{2}]/\mathbb{Z}$ is the quasicyclic group $\mathbb{Z}(2^\infty)$, which is not residually finite.

Definition 2.2. A group G is called *conjugacy separable (CS)* if every conjugacy class $[g]$ is separable.

In a conjugacy separable group G , if two elements $g, h \in G$ are non-conjugate, then their images \bar{g}, \bar{h} are non-conjugate in some finite quotient \bar{G} . Many examples of conjugacy separable groups are known, among which all polycyclic-by-finite groups [6, 18]. Unfortunately, this property behaves quite badly: it is neither subgroup-closed [14, Thm. 1.1], nor quotient-closed (again, $\mathbb{Z}[\frac{1}{2}]$ is a counterexample), nor closed under taking finite extensions [9].

Definition 2.3. A group G is called *twisted conjugacy separable (TCS)* if for every automorphism $\varphi \in \text{Aut}(G)$ and every element $g \in G$, the twisted conjugacy class $[g]_\varphi$ is separable.

This definition was given by Deré and Pengitore in [2]. Little is known about how this property behaves with respect to subgroups, quotients and finite extensions, hence we will study this property extensively in Section 4.

The original definition of twisted conjugacy separability, which differs from Definition 2.3, was given by Fel'shtyn and Troitsky in [5] and further developed in [4]. Their definition says that a group G is twisted conjugacy separable if for every $\varphi \in \text{Aut}(G)$ with finitely many twisted conjugacy classes and for every $g, h \in G$ with $[g]_\varphi \neq [h]_\varphi$, there exists a φ -invariant $N \trianglelefteq_f G$ such that $[\bar{g}]_{\bar{\varphi}} \neq [\bar{h}]_{\bar{\varphi}}$, where $\bar{\varphi}$ is the induced automorphism on $\bar{G} := G/N$. In the sequel, we will exclusively use Definition 2.3.

Definition 2.4. A group G is called *strongly residually finite (SRF)* if every normal subgroup of G is separable.

Equivalently, G is an SRF-group if and only if every quotient of G is residually finite. These groups also go by “QRF-groups” and “(RF)^Q-groups”. Strong residual finiteness was studied in FC-groups [12] and nilpotent groups [15]; any finitely generated abelian-by-polycyclic-by-finite group is SRF [10, Thm. 3]. The family of SRF-groups is closed under taking finite extensions, but in contrast with RF-groups, quotient-closed yet not subgroup-closed. For example, the Baumslag-Solitar group $\text{BS}(1, 2)$ is SRF, but its derived subgroup $\mathbb{Z}[\frac{1}{2}]$ is not.

Definition 2.5. A group G is called *extended residually finite (ERF)* if every subgroup of G is separable.

We refer to [20] for a summary on ERF-groups. The family of ERF-groups is closed under taking finite extensions and is both subgroup- and quotient-closed. A well-known result of Mal'cev says that polycyclic-by-finite groups are ERF [13]; they remain the only known finitely generated ERF-groups. Characterisations for nilpotent [15], FC [20], FC* [21] and linear [27] ERF-groups are known.

In [24], Stebe introduced *nests* and *residual finiteness with respect to nests*, specifically with the aim of uniting conjugacy separability and extended residual finiteness in a single separability property.

Definition 2.6. Let G be a group. A *pre-nest* \mathcal{N}_\times is a non-empty subset of $G \times G$ such that if both $(a, b) \in \mathcal{N}_\times$ and $(c, d) \in \mathcal{N}_\times$, then also $(ac^{-1}, d^{-1}b) \in \mathcal{N}_\times$. A *nest* \mathcal{N} in G is the image of a pre-nest \mathcal{N}_\times in $G \times G$ under the multiplication map $G \times G \rightarrow G: (g_1, g_2) \mapsto g_1g_2$.

It can easily be seen that any pre-nest must contain $(1, 1)$ and hence any nest contains the identity. We also remark that Stebe did not use the term “pre-nest” and instead called these sets “nests” as well. Below are some examples of nests, all of which appeared in [24, 25].

Example 2.7. *The following subsets of a group G are nests:*

- (1) *any subgroup $H \leq G$;*
- (2) *any double coset HK for subgroups $H, K \leq G$;*
- (3) *the set of commutators $\{[h, g] \mid h \in G\}$ for any fixed $g \in G$;*
- (4) *the set $\{h\varphi(h)^{-1} \mid h \in G\}$ for any fixed $\varphi \in \text{Aut}(G)$.*

Definition 2.8. A group G is called *residually finite with respect to nests (NRF)* if every nest in G is separable.

The family of NRF-groups is closed under taking subgroups [24, Thm. 4] and under taking finite extensions [25, Thm. 3]; the behaviour with respect to quotients will be discussed in Section 5.

Example 2.7 (1) shows that being NRF implies being extended residually finite. From (2), we can deduce that in NRF-groups any double coset HgK is separable, since $x \in HgK$ if and only if $xg^{-1} \in HgKg^{-1}$. Similarly, (3) implies that an NRF-group is conjugacy separable, as $x = hgh^{-1}$ if and only if $xg^{-1} = [h, g]$. Finally, (4) implies that an NRF-group is twisted conjugacy separable: $x = hg\varphi(h)^{-1}$ if and only if $xg^{-1} = h(\iota_g\varphi)(h)^{-1}$.

We summarise how these notions of separability behave with respect to taking subgroups, quotients and finite extensions in Table 1. Their relation to one another is illustrated by the diagram below.




TABLE 1. Separability properties of groups

separability property	subgroup-closed	quotient-closed	closed under finite extensions
RF	Yes	No	Yes
CS	No	No	No
TCS	?	?	?
SRF	No	Yes	Yes
ERF	Yes	Yes	Yes
NRF	Yes	?	Yes

Proposition 2.9. *In Diagram (1), the two inclusions in the middle and the two inclusions to the right are indeed strict.*

Proof. We give examples of groups that illustrate that these inclusions are strict.

- $\text{CS} \subsetneq \text{RF}$: the general and special linear groups $\text{GL}_n(\mathbb{Z}), \text{SL}_n(\mathbb{Z})$ with $n \geq 3$ are RF but not CS [23, Thm. 3].
- $\text{TCS} \subsetneq \text{CS}$: if G is a centreless finite group, the direct sum of countably infinitely many copies of G is CS but not TCS [26, Sec. 3].
- $\text{SRF} \subsetneq \text{RF}$: the group of dyadic rationals $\mathbb{Z}[\frac{1}{2}]$ is RF, but its quasicyclic quotient $\mathbb{Z}[\frac{1}{2}]/\mathbb{Z}$ is not. Thus $\mathbb{Z}[\frac{1}{2}]$ is not SRF.
- $\text{ERF} \subsetneq \text{SRF}$: we stated earlier that $\text{BS}(1, 2)$ is SRF, but its derived subgroup $\mathbb{Z}[\frac{1}{2}]$ is not. If $\text{BS}(1, 2)$ were ERF, then its derived subgroup would also be ERF and therefore SRF. \square

3. PROFINITE GROUPS

The class of profinite groups will be a rich source of (counter)examples when studying twisted conjugacy separability. Below, we give a short overview of the definitions and properties we will need later, based on [19, Sec. 2.1 & Sec. 4.2].

Definition 3.1. A *profinite group* is a topological group that is compact, Hausdorff and totally disconnected.

Proposition 3.2. For any profinite group, one has:

- The identity has a local basis consisting of open normal subgroups;
- Subgroups are open if and only if they are closed and have finite index.

Let G be a group and consider the collection of its finite index normal subgroups. If $M, N \trianglelefteq_f G$ and $M \leq N$, let $p_{MN}: G/M \rightarrow G/N$ be the natural projection. Then $\{G/N, p_{MN}\}$ is an inverse system of (finite) groups.

Definition 3.3. The *profinite completion* \hat{G} of a group G is the projective limit

$$\hat{G} := \varprojlim_{N \trianglelefteq_f G} G/N,$$

equipped with the limit topology.

Thus, the profinite completion \hat{G} of G is a subgroup of the direct product $\prod_{N \trianglelefteq_f G} G/N$, and the map

$$\iota: G \rightarrow \hat{G} \subseteq \prod_{N \trianglelefteq_f G} G/N: g \mapsto (gN)_N$$

provides a canonical inclusion of G into \hat{G} .

Proposition 3.4. Let G be a group, \hat{G} its profinite completion and $\iota: G \rightarrow \hat{G}$ the canonical inclusion. Then:

- $\iota(G)$ is dense in \hat{G} ;
- ι is injective if and only if G is residually finite.

Let G be a group and τ_{PF} a topology such that G equipped with τ_{PF} is a profinite group. Let U be an open set in G with respect to τ_{PF} . For any $g \in U$, the set $g^{-1}U$ is open and contains the identity 1_G . Thus, there exists an open normal subgroup N_g contained in $g^{-1}U$. But then

$$U = \bigcup_{g \in U} gN_g,$$

and each N_g has finite index G . Then U is also open in the finite-index topology τ_{FI} on G . Thus, in general, the finite-index topology τ_{FI} on G is finer than τ_{PF} . Those profinite groups for which both topologies coincide, which happens exactly when all finite index subgroups are open, are of particular interest to us.

Definition 3.5. A profinite group G is called *strongly complete* if it satisfies any of the following equivalent conditions:

- (1) Every finite index subgroup of G is open;
- (2) G coincides with its profinite completion \hat{G} ;
- (3) Every homomorphism from G to a profinite group is continuous.

We say that a topological group G is *topologically finitely generated* if it admits a finitely generated subgroup $\langle g_1, \dots, g_n \rangle$ that is dense. In [17], Nikolov and Segal proved the following celebrated theorem.

Theorem 3.6. *Topologically finitely generated profinite groups are strongly complete.*

If G is a finitely generated group and \hat{G} is its profinite completion, then $\iota(G)$ is finitely generated and dense in \hat{G} . Thus, \hat{G} is topologically finitely generated and therefore strongly complete.

4. TWISTED CONJUGACY SEPARABILITY

The definition of twisted conjugacy separability was already given in Section 2. We shall start this section by giving some related definitions, and then study how twisted conjugacy separability behaves with respect to taking subgroups, quotients and finite extensions.

Definition 4.1. A group G is (φ, ψ) -twisted conjugacy separable, where $\varphi, \psi \in \text{Hom}(H, G)$ for some group H , if every (φ, ψ) -twisted conjugacy class is separable.

Equivalently, for any pair g, k of non- (φ, ψ) -twisted conjugate elements of G , there exists a finite quotient \bar{G} such that \bar{g}, \bar{k} are non- $(\bar{\varphi}, \bar{\psi})$ -twisted conjugated, where $\bar{\varphi}, \bar{\psi} \in \text{Hom}(H, \bar{G})$.

Definition 4.2. Let G and H be groups. We say that G is H -twisted conjugacy separable if G is (φ, ψ) -twisted conjugacy separable for all $\varphi, \psi \in \text{Hom}(H, G)$.

We are now ready to study twisted conjugacy separability on profinite groups. The following lemma can be found in e.g. [3, Lem. 2.3(1)].

Lemma 4.3. *Let G and H be profinite groups and let $\varphi, \psi \in \text{Hom}(H, G)$ be continuous homomorphisms. Then the (φ, ψ) -twisted conjugacy classes are closed.*

Proof. Let $g \in G$. The map $H \rightarrow G: h \mapsto \psi(h)g\varphi(h)^{-1}$ is continuous, and its image is exactly $[g]_{\varphi, \psi}$. Because H is compact, so is its continuous image $[g]_{\varphi, \psi}$. Since G is Hausdorff, $[g]_{\varphi, \psi}$ is then closed. \square

Theorem 4.4. *Let G and H be profinite groups, with H being strongly complete. Then G is H -twisted conjugacy separable.*

Proof. Let τ_{PF} be a topology on G such that G equipped with τ_{PF} is profinite. Let $\varphi, \psi \in \text{Hom}(H, G)$, which are necessarily continuous since H is strongly complete. By Lemma 4.3, all (φ, ψ) -twisted conjugacy classes are closed with respect to τ_{PF} . But τ_{PF} is coarser than the finite-index topology τ_{FI} on G . Thus, the twisted conjugacy classes are also closed with respect to τ_{FI} and are therefore separable. \square

When considering automorphisms of a strongly complete profinite group G , we thus get the next result.

Corollary 4.5. *Strongly complete profinite groups are twisted conjugacy separable.*

We now fill in some of the gaps left in Section 2. First, we aim to show that a subgroup of a TCS-group need not be TCS.

Example 4.6. *Let G be a finitely generated RF-group that is not conjugacy separable, e.g. $\text{GL}_n(\mathbb{Z})$ with $n \geq 3$. Let \hat{G} be its profinite completion and $\iota: G \rightarrow \hat{G}$ the canonical inclusion. Then \hat{G} is strongly complete and therefore TCS, but its subgroup $\iota(G) \cong G$ is not even conjugacy separable.*

Similarly, we illustrate how the quotient of a TCS-group need not be TCS.

Example 4.7. Let p be a prime and consider the group of p -adic integers \mathbb{Z}_p , which is a topologically finitely generated profinite group. Thus, it is strongly complete and hence TCS. Let \mathbb{Q}_p denote the p -adic rationals, i.e.

$$\mathbb{Q}_p = \bigcup_{n \in \mathbb{N}} \frac{1}{p^n} \mathbb{Z}_p,$$

which is a vector space over \mathbb{Q} . Using the axiom of choice, pick a \mathbb{Q} -vector space epimorphism $\mathbb{Q}_p \rightarrow \mathbb{Q}$. Its restriction to \mathbb{Z}_p is still surjective (see [16, Sec. 1.2.1]), so \mathbb{Z}_p has a quotient isomorphic to \mathbb{Q} . But \mathbb{Q} is not even residually finite.

Examples 4.6 and 4.7 also show that TCS-groups need not be NRF, since the family of NRF-groups is closed under taking subgroups and quotients. Thus, the inclusion $\text{NRF} \subsetneq \text{TCS}$ is strict.

The final question we would like to answer, is whether or not a finite extension of a TCS-group is again TCS. We provide a partial answer, which is strongly based on [4, Thm. 5.2], but the general case will remain open.

Proposition 4.8. Let N be a finite index normal subgroup of a group G .

- (a) If N is TCS, then G is CS;
- (b) If N is characteristic and TCS, then G is TCS.

Proof. Let $g \in G$. Let x_1, \dots, x_r be coset representatives of G/N with $r := [G : N]$. It is easy to see that if a set is open (closed) in the finite-index topology on H , then it is open (closed) in the finite-index topology on G .

For each $i = 1, \dots, r$, set $g_i := x_i g x_i^{-1}$ and let $\varphi_i \in \text{Aut}(N)$ be the restriction of $\iota_{g_i} \in \text{Inn}(G)$ to N . We can write

$$\begin{aligned} [g] &= \{hgh^{-1} \mid h \in G\} \\ &= \bigcup_{i=1}^r \{nx_i g x_i^{-1} n^{-1} \mid n \in N\} \\ &= \bigcup_{i=1}^r \{ng_i n^{-1} \mid n \in N\} \\ &= \bigcup_{i=1}^r \{n\varphi_i(n)^{-1} g_i \mid n \in N\} \\ &= \bigcup_{i=1}^r [1]_{\varphi_i} \cdot g_i. \end{aligned}$$

For each i , $[1]_{\varphi_i}$ is a twisted conjugacy class in N . Thus, each $[1]_{\varphi_i}$ is closed in N and therefore closed in G . So $[g]$ is a finite union of closed sets and therefore closed itself, which proves part (a).

For part (b), one may take an analogous approach. Let $g \in G$ and $\psi \in \text{Aut}(G)$, set $g_i := x_i g \psi(x_i)^{-1}$ and let $\psi_i \in \text{Aut}(N)$ be the restriction of $\iota_{g_i} \psi$ to N . We can then write

$$[g]_{\psi} = \bigcup_{i=1}^r [1]_{\psi_i} \cdot g_i,$$

so indeed $[g]_{\psi}$ is closed in the finite-index topology on G . \square

Question 4.9. Does there exist a group G with finite index normal subgroup N such that N is TCS but G is not?

5. NESTS AND COMPLETE TWISTED CONJUGACY SEPARABILITY

The main result obtained by Stebe with regards to nests, is that polycyclic-by-finite groups are residually finite with respect to nests [25, Thm. 5]. This result was used by Gonçalves and Wong to study twisted conjugacy. Using the constructions and arguments they provide in [8, Sec. 3], one may extract the following theorem and proof.

Theorem 5.1. *Let G and H be polycyclic-by-finite groups. Then G is H -twisted conjugacy separable.*

Proof. Let $\varphi, \psi \in \text{Hom}(H, G)$ and let $g_1, g_2 \in G$ with $[g_1]_{\varphi, \psi} \neq [g_2]_{\varphi, \psi}$. Consider the subgroups

$$A := \{(h, \psi(h)) \mid h \in H\}, \quad B := \{(h, g_2\varphi(h)g_2^{-1}) \mid h \in H\}$$

of $H \times G$. The double coset AB is a nest in $H \times G$. Suppose, by contradiction, that $(1, g_1g_2^{-1}) \in AB$. Then there exist $h_1, h_2 \in H$ such that

$$(1, g_1g_2^{-1}) = (h_1h_2, \psi(h_1)g_2\varphi(h_2)g_2^{-1}).$$

Looking at the first component, we must have $h_2 = h_1^{-1}$. Equality for the second component then becomes

$$g_1g_2^{-1} = \psi(h_1)g_2\varphi(h_1)^{-1}g_2^{-1}.$$

If such h_1 exists, then g_1 and g_2 would be (φ, ψ) -twisted conjugate, which is untrue by assumption. So $(1, g_1g_2^{-1}) \notin AB$. Let $N \trianglelefteq_f H \times G$ such that

$$(1, g_1g_2^{-1}) \notin ABN.$$

Then, setting $M := G \cap N \trianglelefteq_f G$, we find that $g_1 \notin [g_2]_{\varphi, \psi}M$. \square

However, there is an alternative way to apply the separability of nests to (φ, ψ) -twisted conjugacy classes.

Proposition 5.2. *Let G and H be groups and let $\varphi, \psi \in \text{Hom}(H, G)$. Then the twisted conjugacy class $[1]_{\varphi, \psi}$ is a nest in G .*

Proof. The set $\{(\psi(h), \varphi(h)^{-1}) \mid h \in H\}$ is a pre-nest in $G \times G$. Its corresponding nest in G is exactly $[1]_{\varphi, \psi}$. \square

At first glance, this result does not appear to be all that useful. After all, we want to study separability of *all* twisted conjugacy classes, and not just the class of the identity. The following lemma and proposition show that studying the separability of the twisted conjugacy class of the identity is, in fact, sufficient.

Lemma 5.3. *Let G and H be groups, let $g, k \in G$ and let $\varphi, \psi \in \text{Hom}(H, G)$. Then*

$$[g]_{\varphi, \psi} = [k]_{\varphi, \psi} \iff [gk^{-1}]_{\iota_k \varphi, \psi} = [1]_{\iota_k \varphi, \psi}.$$

Proposition 5.4. *Let G and H be groups. The following are equivalent:*

- (1) G is H -twisted conjugacy separable.
- (2) For all $\varphi, \psi \in \text{Hom}(H, G)$, the twisted conjugacy class $[1]_{\varphi, \psi}$ is separable.

Proof. It is easy to see that (1) implies (2). For the reverse implication, let $g, k \in G$ and $\varphi, \psi \in \text{Hom}(H, G)$ with $[g]_{\varphi, \psi} \neq [k]_{\varphi, \psi}$. Applying Lemma 5.3, we find that $[gk^{-1}]_{\iota_k \varphi, \psi} \neq [1]_{\iota_k \varphi, \psi}$. Then there exists $N \trianglelefteq_f G$ such that $[\bar{g}k^{-1}]_{\iota_{\bar{k}} \bar{\varphi}, \bar{\psi}} \neq [\bar{1}]_{\iota_{\bar{k}} \bar{\varphi}, \bar{\psi}}$ in $\bar{G} := G/N$, where $\bar{\varphi}, \bar{\psi} \in \text{Hom}(H, \bar{G})$. Applying Lemma 5.3 again, we get $[\bar{g}]_{\bar{\varphi}, \bar{\psi}} \neq [\bar{k}]_{\bar{\varphi}, \bar{\psi}}$. \square

Since polycyclic-by-finite groups are residually finite with respect to nests, and the twisted conjugacy class $[1]_{\varphi, \psi}$ is always a nest irrespective of the group H and the homomorphisms $\varphi, \psi: H \rightarrow G$, Proposition 5.4 implies that polycyclic-by-finite groups are H -twisted conjugacy separable for every group H . We give this property a name:

Definition 5.5. A group G is *completely twisted conjugacy separable (CTCS)* if it is H -twisted conjugacy separable for every group H .

At this point, it is clear that being residually finite with respect to nests implies complete twisted conjugacy separability. To prove the converse, we first need to understand the structure of pre-nests, which was described by Stebe in [24, Thm. 1].

Theorem 5.6. *Let S_1 and S_2 be subgroups of a group G . Let K_i be a normal subgroup of S_i for $i = 1, 2$ such that S_1/K_1 is isomorphic to S_2/K_2 , and let θ be an isomorphism between the factor groups. The set of all elements of $G \times G$ of the form $(ac, \theta(c^{-1})b)$, with a in K_1 , b in K_2 and c a coset representative of S_1 modulo K_1 is a pre-nest in $G \times G$. Conversely, every pre-nest in $G \times G$ may be obtained by this construction.*

It should be noted that there is some slight abuse of notation in the theorem above: “ $\theta(c^{-1})$ ” should be interpreted as “a coset representative of $\theta(c^{-1}K_1)$ ”. In the case of a (φ, ψ) -twisted conjugacy class with $\varphi, \psi \in \text{Hom}(H, G)$, the groups in question are given by

$$\begin{aligned} S_1 &= \psi(H) & K_1 &= \psi(\ker(\varphi)) \\ S_2 &= \varphi(H) & K_2 &= \varphi(\ker(\psi)) \end{aligned}$$

and θ is given by

$$\theta: \frac{S_1}{K_1} \rightarrow \frac{S_2}{K_2}: \psi(h)K_1 \mapsto \varphi(h)K_2.$$

Using the structure description of pre-nests, we may prove the following converse of Proposition 5.2.

Proposition 5.7. *Let G be a group and \mathcal{N} a nest in G . Then there exist $H \leq G \times G$ and $\varphi, \psi \in \text{Hom}(H, G)$ such that $\mathcal{N} = [1]_{\varphi, \psi}$.*

Proof. Let \mathcal{N} be a nest in G and let \mathcal{N}_\times be a pre-nest in $G \times G$ inducing \mathcal{N} . Let $S_1, S_2, K_1, K_2, \theta$ determine \mathcal{N}_\times as in Theorem 5.6. Consider the subgroup $D \leq S_1/K_1 \times S_2/K_2$ given by

$$D := \{(\bar{c}, \theta(\bar{c})) \mid \bar{c} \in S_1/K_1\}.$$

For every $\bar{c} \in S_1/K_1$, fix some preimage $c \in S_1$. Similarly, with the same abuse of notation as mentioned before, fix a preimage $\theta(c) \in S_2$ of $\theta(\bar{c})$. Then H , the preimage of D in $S_1 \times S_2$, is exactly the set

$$H = \{(ac, b^{-1}\theta(c)) \mid a \in K_1, b \in K_2, \bar{c} \in S_1/K_1\}.$$

Let ψ and φ be the projections to the first and second component respectively, composed with the inclusion in G , such that $\psi, \varphi \in \text{Hom}(H, G)$. It is now easy to see that

$$\mathcal{N}_\times = \{(\psi(h), \varphi(h)^{-1}) \mid h \in H\},$$

and therefore $\mathcal{N} = [1]_{\varphi, \psi}$. □

Combined, Propositions 5.2 and 5.7 lead us to our first main result:

Theorem A. *A group is residually finite with respect to nests if and only if it is completely twisted conjugacy separable.*

Stebe already showed that the family of NRF-groups is closed under taking subgroups and finite extensions, but he did not study quotients. We do so below.

Proposition 5.8. *The family of NRF-groups is closed under taking quotients.*

Proof. Let G be an NRF-group, $N \trianglelefteq G$, and let p be the natural projection from G to $\bar{G} := G/N$. Suppose that $\bar{g} \in \bar{G}$ and that $\bar{\mathcal{N}}$ is a nest in \bar{G} with $\bar{g} \notin \bar{\mathcal{N}}$. Let $\bar{\mathcal{N}}_\times$ be a pre-nest in $\bar{G} \times \bar{G}$ corresponding to $\bar{\mathcal{N}}$. Then the set

$$\mathcal{N}_\times := \{(g_1, g_2) \in G \mid (p(g_1), p(g_2)) \in \bar{\mathcal{N}}_\times\}$$

is a pre-nest in $G \times G$, and the corresponding nest \mathcal{N} in G is exactly $p^{-1}(\bar{\mathcal{N}})$. Let $g \in p^{-1}(\bar{g})$, then $g \notin \mathcal{N}$ and thus there exists some $K \trianglelefteq_f G$ such that $g \notin \mathcal{N}K$. Set $\bar{K} := p(K)$.

Suppose, by contradiction, that $\bar{g} \in \bar{\mathcal{N}}\bar{K}$. Then there exist $n \in N$, $h \in \mathcal{N}$ and $k \in K$ such that $g = nhk$. But $nh \in \mathcal{N}$, so $g \in \mathcal{N}K$, which contradicts the choice of K . Therefore indeed $\bar{g} \notin \bar{\mathcal{N}}\bar{K}$ with $\bar{K} \trianglelefteq_f \bar{G}$. \square

We can update Table 1 and Diagram (1) with our new findings, the results can be found in Table 2 and Diagram (2).

$$(2) \quad \begin{array}{ccc} \text{NRF} \subseteq \text{ERF} \subsetneq \text{SRF} & & \\ \parallel & & \curvearrowright \text{RF} \\ \text{CTCS} \subsetneq \text{TCS} \subsetneq \text{CS} & & \curvearrowleft \end{array}$$

TABLE 2. Separability properties of groups (revisited)

separability property	subgroup-closed	quotient-closed	closed under finite extensions
RF	Yes	No	Yes
CS	No	No	No
TCS	No	No	?
CTCS	Yes	Yes	Yes
SRF	No	Yes	Yes
ERF	Yes	Yes	Yes
NRF	Yes	Yes	Yes

In Diagram (2), one uncertainty remains: whether the inclusion $\text{ERF} \subseteq \text{NRF}$ is strict or an equality.

Question 5.9. Does there exist an ERF-group that is not NRF?

6. NILPOTENT-BY-FINITE GROUPS

The study of twisted conjugacy on nilpotent-by-finite groups has proven to be very successful. As such, this family of groups is natural starting point to study complete twisted conjugacy separability. Since finitely generated nilpotent-by-finite groups are polycyclic-by-finite, we already know they are completely twisted conjugacy separable.

Thus, we consider a slightly larger family of groups, being the nilpotent-by-finite groups that are strongly residually finite. These groups were characterised by Menth [15, Thm. 5.15], see also [20, Sec. 4] and [27, Sec. 2] for more details and remarks.

Theorem 6.1. *Let G be a nilpotent group with torsion subgroup $\tau(G)$. Then G is SRF if and only if the following conditions hold:*

- $G/\tau(G)$ has no quasicyclic section;
- each p -component of $\tau(G)$ is abelian-by-finite and has finite exponent.

For the reader's convenience, we provide an example of an infinitely generated nilpotent SRF-group, which appeared in both [21, Ex. 4.1] and [27, Lem. 2.8].

Example 6.2. *The group $G := (\bigoplus_p C_{p^2}) \rtimes \mathbb{Z}$, where $\mathbb{Z} = \langle x \rangle$ acts on the cyclic group $C_{p^2} = \langle y_p \rangle$ via $y_p^x = y_p^{1+p}$ for each prime p , is a 2-step nilpotent SRF-group.*

Smirnov proved in [22] that for nilpotent groups, the properties SRF and ERF are equivalent; Menth generalised this in [15, Thm. 7.3] to nilpotent-by-finite groups. We aim to extend these results to complete twisted conjugacy separability, for which we will need some intermediate results.

The lemma below is inspired by [3, Lem. 2.7] and [20, Cor. 4.3].

Lemma 6.3. *Let \mathcal{P} be a property that is retained when taking subgroups and finite extensions. Then a finite-by- \mathcal{P} RF-group also has \mathcal{P} .*

Proof. Let G be residually finite and let $N \trianglelefteq G$ be a finite subgroup such that G/N has \mathcal{P} . Take $M \trianglelefteq_f G$ such that $M \cap N = 1$. By the second isomorphism theorem,

$$M \cong \frac{M}{M \cap N} \cong \frac{MN}{N} \leq \frac{G}{N},$$

hence M has \mathcal{P} . Then G has \mathcal{P} as well. \square

Proposition 6.4. *Let G be an SRF-group. If C is a central subgroup of G such that G/C is CTCS, then G is CTCS.*

Proof. Let $g \in G$ and $\varphi, \psi \in \text{Hom}(H, G)$ with $[g]_{\varphi, \psi} \neq [1]_{\varphi, \psi}$. Set $\tilde{G} := G/C$. We consider two cases.

First, if $[\tilde{g}]_{\tilde{\varphi}, \tilde{\psi}} \neq [\tilde{1}]_{\tilde{\varphi}, \tilde{\psi}}$, then we find $\tilde{N} \trianglelefteq_f \tilde{G}$ such that $[\tilde{g}]_{\tilde{\varphi}, \tilde{\psi}} \neq [\tilde{1}]_{\tilde{\varphi}, \tilde{\psi}}$ in $\tilde{G} := \tilde{G}/\tilde{N}$. Let N be the preimage of \tilde{N} in G , then $N \trianglelefteq_f G$ and hence $\tilde{G} \cong G/N$ is a finite quotient of G .

Second, if $[\tilde{g}]_{\tilde{\varphi}, \tilde{\psi}} = [\tilde{1}]_{\tilde{\varphi}, \tilde{\psi}}$, we have that $[g]_{\varphi, \psi} = [c]_{\varphi, \psi}$ for some $c \in C$. Set $D := C \cap [1]_{\varphi, \psi}$, which is a subgroup because C is central. As G is SRF and D is normal in G , pick some $M \trianglelefteq_f G$ such that $c \notin DM$. Set $K := M \cap C$, then K is normal in G , $K \trianglelefteq_f C$ and $c \notin DK$. Set $\tilde{G} := G/K$ and suppose, by contradiction, that $[\tilde{c}]_{\tilde{\varphi}, \tilde{\psi}} = [\tilde{1}]_{\tilde{\varphi}, \tilde{\psi}}$. Then there exist $h \in G$ and $k \in K$ such that

$$c = \psi(h)\varphi(h)^{-1}k.$$

But then $\psi(h)\varphi(h)^{-1} = ck^{-1} \in C$, and hence $\psi(h)\varphi(h)^{-1} \in D$. In turn, this means that $c \in DK$, which cannot be the case. Therefore $[\tilde{c}]_{\tilde{\varphi}, \tilde{\psi}} \neq [\tilde{1}]_{\tilde{\varphi}, \tilde{\psi}}$. Now \tilde{G} is finite-by-CTCS and residually finite. Using Lemma 6.3, it is CTCS. We can now proceed as in the first case. \square

We now have all necessary tools to extend Menth's result to complete twisted conjugacy separability.

Theorem 6.5. *For a nilpotent-by-finite group G , the following are equivalent:*

- (a) G is strongly residually finite (SRF);
- (b) G is extended residually finite (ERF);
- (c) G is residually finite with respect to nests (NRF);
- (d) G is completely twisted conjugacy separable (CTCS).

Proof. The equivalence between (a) and (b) is the aforementioned result by Menth, and the equivalence between (c) and (d) is just Theorem A.

Since any subgroup is a nest, (c) implies (b), and it now suffices to prove that (b) implies (d). Let G be a nilpotent-by-finite ERF-group and let $N \trianglelefteq_f G$ be nilpotent. Because subgroups of ERF-groups are ERF, N is ERF and hence SRF. By induction on the nilpotency class and using Proposition 6.4, it follows that N is CTCS. Since G is a finite extension of N , G itself is CTCS. So indeed (b) implies (d) and hence all of (a), (b), (c) and (d) are equivalent. \square

7. POLYCYCLIC-BY-NILPOTENT-BY-FINITE GROUPS

At this point, we know of two families of groups that are completely twisted conjugacy separable: the polycyclic-by-finite groups and the nilpotent-by-finite SRF-groups, which are described in Theorem 6.1. Using a result by Kilsch, we can glue these results together and extend Theorem 6.5 to the family of polycyclic-by-nilpotent-by-finite groups.

A priori, the ‘‘polycyclic-by-nilpotent-by-finite’’ could be ambiguous, as it could refer to either polycyclic-by-(nilpotent-by-finite) groups or (polycyclic-by-nilpotent)-by-finite groups. The following proposition shows that there is no such ambiguity.

Proposition 7.1. *Let G be a group. Then G is polycyclic-by-(nilpotent-by-finite) if and only if it is (polycyclic-by-nilpotent)-by-finite.*

Proof. First assume that G is polycyclic-by-(nilpotent-by-finite). Let $P \trianglelefteq G$ be polycyclic normal subgroup such that G/P is nilpotent-by-finite. Then there exists a nilpotent finite index normal subgroup of G/P , which must be of the form N/P . This N is a finite index normal subgroup of G that is polycyclic-by-nilpotent, therefore G is (polycyclic-by-nilpotent)-by-finite.

For the converse, let $N \trianglelefteq_f G$ be polycyclic-by-nilpotent and let $P \trianglelefteq N$ be polycyclic such that N/P is nilpotent. If c is the nilpotency class of N/P , then $\gamma_{c+1}(N/P) = 1$. But

$$\gamma_{c+1}(N/P) = \frac{\gamma_{c+1}(N)P}{P},$$

thus $\gamma_{c+1}(N) \leq P$, hence $\gamma_{c+1}(N)$ is then polycyclic. Moreover, since it is characteristic in N , it is normal in G . Therefore $G/\gamma_{c+1}(N)$ contains $N/\gamma_{c+1}(N)$ as a nilpotent finite index normal subgroup, and G is polycyclic-by-(nilpotent-by-finite). \square

We will use a result by Kilsch on derivations. The definition used by Kilsch is that a derivation $d: H \rightarrow G$ between groups H and G is a map such that $d(h_1 h_2) = d(h_1)^{h_2} d(h_2)$, where H acts on G (from the right) via automorphisms.

Theorem 7.2 (see [11, Thm. D]). *The image of a derivation from a soluble-by-finite group into a polycyclic-by-finite group G is closed with respect to the finite-index topology on G .*

Definition 7.3. Let G and H be groups and let $\varphi, \psi \in \text{Hom}(H, G)$. The *coincidence subgroup* $\text{Coin}(\varphi, \psi)$ of φ and ψ is defined as

$$\text{Coin}(\varphi, \psi) := \{h \in H \mid \varphi(h) = \psi(h)\}.$$

We now generalise Theorem 6.5 to the family of polycyclic-by-nilpotent-by-finite groups, obtaining our second main result.

Theorem B. *For a polycyclic-by-nilpotent-by-finite group G , the following are equivalent:*

- (a) G is strongly residually finite (SRF);
- (b) G is extended residually finite (ERF);
- (c) G is residually finite with respect to nests (NRF);
- (d) G is completely twisted conjugacy separable (CTCS).

Proof. It suffices to prove that G being SRF implies it being CTCS. Let $g \in G$, let H be a group and let $\varphi, \psi \in \text{Hom}(H, G)$ such that $[g]_{\varphi, \psi} \neq [1]_{\varphi, \psi}$. We may assume that $H \leq G \times G$ and therefore that H is soluble-by-finite. Let $N \trianglelefteq G$ be a polycyclic normal subgroup such that $\bar{G} := G/N$ is nilpotent-by-finite, and let $\bar{\varphi}, \bar{\psi} \in \text{Hom}(H, \bar{G})$. If $[\bar{g}]_{\bar{\varphi}, \bar{\psi}} \neq [\bar{1}]_{\bar{\varphi}, \bar{\psi}}$, then we need just apply Theorem 6.5.

If however $[\bar{g}]_{\bar{\varphi}, \bar{\psi}} = [\bar{1}]_{\bar{\varphi}, \bar{\psi}}$, then there exists some $n \in N$ such that $[g]_{\varphi, \psi} = [n]_{\varphi, \psi}$. Now consider the map

$$d: \text{Coin}(\bar{\varphi}, \bar{\psi}) \rightarrow N: h \mapsto \psi(h)^{-1} \varphi(h).$$

For $h_1, h_2 \in \text{Coin}(\bar{\varphi}, \bar{\psi})$, one has that

$$\begin{aligned} d(h_1 h_2) &= \psi(h_1 h_2)^{-1} \varphi(h_1 h_2) \\ &= \psi(h_2)^{-1} \psi(h_1)^{-1} \varphi(h_1) \varphi(h_2) \\ &= \psi(h_2)^{-1} d(h_1) \psi(h_2) \psi(h_2)^{-1} \varphi(h_2) \\ &= d(h_1)^{h_2} d(h_1), \end{aligned}$$

where $\text{Coin}(\bar{\varphi}, \bar{\psi})$ acts on N (from the right) via

$$\text{Coin}(\bar{\varphi}, \bar{\psi}) \rightarrow \text{Aut}(N): h \mapsto \iota_{\psi(h)^{-1}}.$$

Thus, d is a derivation, and its image is exactly $N \cap [1]_{\varphi, \psi}$. Since N is polycyclic and $\text{Coin}(\bar{\varphi}, \bar{\psi}) \leq H$ is soluble-by-finite, we can apply Theorem 7.2: choose $K \trianglelefteq_f N$ such that $n \notin (N \cap [1]_{\varphi, \psi})K$. Since N is finitely generated, we may assume K is characteristic in N and therefore normal in G . Set $\tilde{N} := N/K$ and $\tilde{G} := G/K$, then $[\tilde{g}]_{\tilde{\varphi}, \tilde{\psi}} = [\tilde{n}]_{\tilde{\varphi}, \tilde{\psi}} \neq [\tilde{1}]_{\tilde{\varphi}, \tilde{\psi}}$.

As \tilde{N} is finite and $\tilde{G}/\tilde{N} \cong G/N = \bar{G}$ is nilpotent-by-finite, \tilde{G} is a finite-by-(nilpotent-by-finite) SRF-group. From Lemma 6.3 we obtain that \tilde{G} is then nilpotent-by-finite, hence we can finish by again applying Theorem 6.5. \square

In [13, Thm. 4], Mal'cev proved that the direct product of two ERF-groups is itself ERF, and in [1, Thm. 4] Allenby and Gregorac proved that the semidirect product $N \rtimes H$ of a finitely generated ERF-group N and an ERF-group H is itself ERF. Combined with Theorem B, these results can be used to construct polycyclic-by-nilpotent-by-finite CTCS-groups that are neither polycyclic-by-finite nor nilpotent-by-finite.

Corollary 7.4. *Let $G := N \rtimes H$ be the (semi)direct product of a polycyclic group N and a nilpotent-by-finite ERF-group H . Then G is completely twisted conjugacy separable.*

We now use this corollary to construct some explicit examples.

Example 7.5. *Let N be the group $\mathbb{Z}^2 \rtimes_{\theta} \mathbb{Z}$ with $\theta: \mathbb{Z} \rightarrow \text{Aut}(\mathbb{Z}^2) \cong \text{GL}_2(\mathbb{Z})$ determined by*

$$\theta(1) = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix},$$

which is polycyclic but not nilpotent-by-finite (see e.g. [7, Ex. 3.4]). Let H be the nilpotent group from Example 6.2, which is not polycyclic-by-finite since it is not finitely generated. Then the direct product $N \times H$ is a polycyclic-by-nilpotent CTCS-group.

Example 7.6. Let H again be the nilpotent group from Example 6.2. Consider the homomorphism $\lambda: H \rightarrow \text{Aut}(\mathbb{Z}^2) \cong \text{GL}_2(\mathbb{Z})$, defined by

$$\lambda(x) = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}, \quad \lambda(y_2) = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \lambda(y_p) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \text{ when } p \geq 3.$$

Then $\mathbb{Z}^2 \rtimes_{\lambda} H$ is a polycyclic-by-nilpotent CTCS-group. But this group contains both N (as above) and H as subgroups, so it is neither nilpotent-by-finite nor polycyclic-by-finite.

ACKNOWLEDGEMENTS

The author is indebted to Peter Wong for bringing the work by Stebe on nests to his attention, and to Karel Dekimpe for catching a mistake in the proof of Theorem B and helping to correct said proof.

REFERENCES

- [1] R. B. J. T. Allenby and R. J. Gregorac. ‘On Locally Extended Residually Finite Groups’. In: *Conference on Group Theory*. Vol. 319. Lecture Notes in Mathematics. Springer, 1973, pp. 9–17. DOI: 10/c9w7nq.
- [2] J. Deré and M. Pengitore. ‘Effective twisted conjugacy separability of nilpotent groups’. In: *Math. Z.* 292.3-4 (2018), pp. 763–790. DOI: 10/grzsvq.
- [3] A. Fel’shtyn and B. Klopsch. ‘Pólya-Carlson dichotomy for coincidence Reidemeister zeta functions via profinite completions’. In: *Indag. Math.* 33.4 (2022), pp. 753–767. DOI: 10/grvprk.
- [4] A. Fel’shtyn and E. Troitsky. ‘Twisted conjugacy separable groups’. 2006. arXiv: math/0606764 [math.GR].
- [5] A. Fel’shtyn and E. Troitsky. ‘Twisted Burnside-Frobenius theory for discrete groups’. In: *J. Reine Angew. Math.* 2007.613 (2007), pp. 193–210. DOI: 10/fh5nsd.
- [6] E. Formanek. ‘Conjugate separability in polycyclic groups’. In: *J. Algebra* 42.1 (1976), pp. 1–10. DOI: 10/fgqcps.
- [7] D. Gonçalves and P. Wong. ‘Twisted Conjugacy Classes in Exponential Growth Groups’. In: *Bull. Lond. Math. Soc.* 35.02 (2003), pp. 261–268. DOI: 10/bccmxp.
- [8] D. Gonçalves and P. Wong. ‘Homogeneous spaces in coincidence theory II’. In: *Forum Math.* 17.2 (2005), pp. 297–313. DOI: 10/bxsvc7.
- [9] A. V. Goryaga. ‘Example of a finite extension of an FAC-group that is not an FAC-group’. In: *Sibirsk. Mat. Zh.* 27.3 (1986), pp. 203–205.
- [10] A. V. Jategaonkar. ‘Integral group rings of polycyclic-by-finite groups’. In: *J. Pure Appl. Algebra* 4.3 (1974), pp. 337–343. DOI: 10/cffb7.
- [11] D. Kilsch. ‘Conjugacy separability and separable orbits’. English. In: *J. Pure Appl. Algebra* 30 (1983), pp. 167–179. DOI: 10/fhr4mf.
- [12] L. A. Kurdachenko and J. Otal. ‘FC-Groups All of Whose Factor Groups Are Residually Finite’. In: *Commun. Algebra* 31.3 (2003), pp. 1235–1251. DOI: 10/dn7d6d.
- [13] A. I. Mal’cev. ‘On homomorphisms onto finite groups’. In: *Ivanov. Gos. Ped. Inst. Učen. Zap.* 18 (1958), pp. 49–60.
- [14] A. Martino and A. Minasyan. ‘Conjugacy in normal subgroups of hyperbolic groups’. In: *Forum Math.* 24.5 (2012), pp. 889–909. DOI: 10/cscfns.
- [15] M. Menth. ‘Nilpotent groups with every quotient residually finite’. In: *J. Group Theory* 5.2 (2002), pp. 199–217. DOI: 10/ftf7q5.
- [16] N. Nikolov and D. Segal. ‘Generators and commutators in finite groups; abstract quotients of compact groups’. In: *Invent. Math.* 190.3 (2012), pp. 513–602. DOI: 10/gkqpwx.

- [17] N. Nikolov and D. Segal. ‘On finitely generated profinite groups, I: strong completeness and uniform bounds’. In: *Ann. Math.* 165.1 (2007), pp. 171–238. DOI: 10/cfkrhb.
- [18] V. N. Remeslennikov. ‘Conjugacy in polycyclic groups’. In: *Algebra Logika* 8.6 (1969), pp. 712–725.
- [19] L. Ribes and P. Zalesskii. *Profinite Groups*. 2nd ed. *Ergeb. Math. Grenzgeb.* 40. Springer, 2010. DOI: 10/b8n6t8.
- [20] D. J. S. Robinson, A. Russo and G. Vincenzi. ‘On groups whose subgroups are closed in the profinite topology’. In: *J. Pure Appl. Algebra* 213.4 (2009), pp. 421–429. DOI: 10/cfvt7m.
- [21] D. J. S. Robinson, A. Russo and G. Vincenzi. ‘On the theory of generalized FC-groups’. In: *J. Algebra* 326.1 (2011), pp. 218–226. DOI: 10/fp72vn.
- [22] D. M. Smirnov. ‘On the theory of finitely approximable groups’. In: *Ukraiń. Mat. Ž.* 15 (1963), pp. 453–457.
- [23] P. F. Stebe. ‘Conjugacy separability of groups of integer matrices’. In: *Proc. Amer. Math. Soc.* 32.1 (1972), pp. 1–7. DOI: 10/dsrbnz.
- [24] P. F. Stebe. ‘Nests in nilpotent groups’. In: *Houston J. Math.* 2.3 (1976), pp. 419–426.
- [25] P. F. Stebe. ‘A residual property of certain linear groups’. In: *Trans. Am. Math. Soc.* 302.1 (1987), pp. 333–340. DOI: 10/cn4dm4.
- [26] E. Troitsky. ‘Two Examples Related to the Twisted Burnside-Frobenius Theory for Infinitely Generated Groups’. In: *J. Math. Sci.* 248.5 (2020), pp. 661–666. DOI: 10/grvprj.
- [27] B. A. F. Wehrfritz. ‘Linear groups with all subgroups profinitely closed’. In: *Q. J. Math.* 62.2 (2011), pp. 501–512. DOI: 10/d2x3pz.

KU LEUVEN KULAK KORTRIJK CAMPUS, E. SABBELAAN 53, 8500 KORTRIJK, BELGIUM
Email address: sam.tertooy@kuleuven.be